


Note: Figure not drawn to scale.

Block A of mass 4.0 kg is on a horizontal, frictionless tabletop and is placed against a spring of negligible mass and spring constant 650 N/m. The other end of the spring is attached to a wall. The block is pushed toward the wall until the spring has been compressed a distance x , as shown above. The block is released and follows the trajectory shown, falling 0.80 m vertically and striking a target on the floor that is a horizontal distance of 1.2 m from the edge of the table. Air resistance is negligible.

- Calculate the time elapsed from the instant block A leaves the table to the instant it strikes the floor.
- Calculate the speed of the block as it leaves the table.
- Calculate the distance x the spring was compressed.

Block B, also of mass 4.0 kg, is now placed at the edge of the table. The spring is again compressed a distance x , and block A is released. As it nears the end of the table, it instantaneously collides with and sticks to block B. The blocks follow the trajectory shown in the figure below and strike the floor at a horizontal distance d from the edge of the table.



Note: Figure not drawn to scale.

(d)

Calculate d if x is equal to the value determined in part (c).

(e) Consider the system consisting of the spring, the blocks, and the table. How does the total mechanical energy E_2 of the system just before the blocks leave the table compare to the total mechanical energy E_1 of the system just before block A is released?

☐ $E_1 > E_2$ ☐ $E_2 = E_1$ ☐ $E_2 > E_1$